

A Note about Torsional Rigidity and Euclidean Moment of Inertia of Plane Domains

R. G. Salakhudinov*

(Submitted by F. G. Avkhadiev)

*N. I. Lobachevskii Institute of Mathematics and Mechanics, Kazan (Volga region) Federal University,
ul. Kremlevskaya 18, Kazan, Tatarstan, 420008 Russia*

Received December 16, 2017

Abstract—Denote by $\mathbf{P}(G)$ the torsional rigidity of a simply connected plane domain G , and by $\mathbf{I}_2(G)$ the Euclidean moment of inertia of G . In 1995 F.G. Avkhadiev proved that $\mathbf{P}(G)$ and $\mathbf{I}_2(G)$ are comparable quantities in sense of Pólya and Szegő. Moreover, it was shown that the ratio $\mathbf{P}(G)/\mathbf{I}_2(G)$ belongs to the segment $[1, 64]$. We investigate the following conjecture $\mathbf{P}(G) \geq 3\mathbf{I}_2(G)$, where G is a simply connected domain. We prove that the conjecture is true for polygonal domains circumscribed about a circle. For convex domains we show sharp isoperimetric inequalities, which justify the conjecture, in particular, we prove that $\mathbf{P}(G) > 2\mathbf{I}_2(G)$. Some aspects of approximate formulas for $\mathbf{P}(G)$ are also discussed.

DOI: 10.1134/S1995080218060161

Keywords and phrases: *Torsional rigidity, Warping function, Euclidean moments of a domain with respect to its boundary, Modified torsional rigidity, Inradius of a domain, Isoperimetric inequality, Convex domain.*

1. INTRODUCTION AND SOME CONJECTURES

In analytical studies of steady field problems of the theory of elasticity [1] and microfluid mechanics [2] a boundary value problem

$$\Delta u = -2 \text{ in } G, \quad u = 0 \text{ on } \partial G \quad (1)$$

is solved for complex-shaped domains G , here ∂G is the boundary curve of G . A classical result of the theory in partial differential equations is that there is a unique solution of the boundary value problem (1) for a simply connected domain G . Isoperimetric estimates in these solutions (see, for instance, [3, 4]) require evaluation of the functional

$$\mathbf{P}(G)^{(1)} := 2 \int_G u(x, G) dA. \quad (2)$$

This functional is called the *torsional rigidity* of G in the theory of elasticity, and the *flow rate* in hydrodynamics of pipe flows.

Investigations of the physically important boundary value problem (1), and the physical functional (2) go back to works of B. de Saint-Venant [5], and Lord Rayleigh (see [3, 6]). Estimates of $\mathbf{P}(G)$, using different geometrical or (and) physical characteristics of G , intensively studied by mathematicians since the middle of XX century, when their interest was stimulated by the famous monograph of G. Pólya and G. Szegő [3]. Then a number of works [6] is growing rapidly in this area. There are a number of isoperimetric inequalities [6] for $\mathbf{P}(G)$ in this branch of mathematical physics.

*E-mail: rsalakhud@gmail.com

¹⁾Throughout the paper we will use the bold face for notations of functionals.